arXiv:1401.2950v1 [math.GR] 10 Jan 2014

On a problem of A. Nagy concerning permutable semigroups satisfying a non-trivial permutation identity[1]

Attila Deák

Communicated by M. B. Szendrei

Abstract

We say that a semigroup S is a permutable semigroup if, for every congruences α and β of S, $\alpha \circ \beta = \beta \circ \alpha$. In [4], A. Nagy showed that every permutable semigroup satisfying an arbitrary non-trivial permutation identity is medial or an ideal extension of a rectangular band by a non-trivial commutative nil semigroup. The author raised the following problem: Is every permutable semigroup satisfying a non-trivial permutation identity medial? In present paper we give a positive answer for this problem.

A semigroup S is called a *permutable semigroup* if the congruences of S commute with each other, that is, $\alpha \circ \beta = \beta \circ \alpha$ is satisfied for every congruences α and β of S.

Theorem 1 *Let S be an ideal extension of a rectangular band K by a non trivial commutative nil semigroup N.*
If S is permutable and satisfies a non-trivial permutation identity then either S is medial or N is nilpotent.

Proof. Assume that a permutable semigroup S satisfies a permutation identity

$$x_1 x_2 \dots x_n = x_{\sigma(1)} x_{\sigma(2)} \dots x_{\sigma(n)}, \ (\sigma \neq id).$$

We can suppose that $n > 2$. Let l and k be non-negative integers such that

$$\sigma(1) = 1, \sigma(2) = 2, \cdots, \sigma(l) = l, \sigma(l+1) \neq l+1$$

and

$$\sigma(n-k) \neq n-k, \sigma(n-k+1) = n-k+1, \cdots, \sigma(n) = n.$$

If $l = k = 0$ then, by Theorem 1 of [4], S is commutative. Consider the case when $l > 0$ or $k > 0$. Let j be $\max\{l, k\}$.

Assume that $j = 1$. Then let β be a relation on S such that $a \,\beta\, b \Leftrightarrow uabv = ubav$ for every $u, v \in S$. It is clear that β is reflexive and symmetric. We show that β is transitive. Let $u, v, a, b, c \in S$ be arbitrary elements such that $a \,\beta\, b$ and $b \,\beta\, c$. If $ac \in S - K$ then $ac = ca$, because N is commutative. Thus $a \,\beta\, c$. Assume that $ac \in K$. Then $ca \in K$ and

$$ac = ac(xac)ac = acxac, \ ca = ca(xca)ca = caxca$$

[1]Research supported by the Hungarian NFSR grant No T042481
AMS Subject classification number: 20M10

for every $x \in S$. Thus

$$uacv = (ua)cb(acv) = uab(cacv) = ubacacv = ubacv = uabcv = uacbv.$$

Similarly,

$$ucav = ubcav = ucbav = ucabv.$$

From these results it follows that

$$uacv = ub^p ab^q cb^r v$$

for every non-negative integers p, q, r. Similarly,

$$ucav = ub^p cb^q ab^r v$$

for every non-negative integers p, q, r. Let $\hat{l},\ \hat{k}$ be such that $\sigma(l+1) = \hat{l},\ \sigma(n-k) = \hat{k}$. It is obvious that $\hat{l} \geq l+2$ and $\hat{k} \leq n-k-1$. So

$$uacv = uacb^{n-2}v = ub^l ab^{\hat{l}-l-2}cb^{n-\hat{l}}v = ub^l cb^t ab^s v = ucav,$$

where s, t are non-negative integers such that $s+t+l = n-2$. Hence β is an equivalence on S. Let $x, y \in K$. We will show that $x \,\beta\, y$. Let $u, v \in S$ be arbitrary elements. Then

$$uxyv = ux^{\hat{l}-l-1}y^{n-\hat{l}}yv = uyxyv = uyv = uy^{\hat{k}}xy^{n-\hat{k}-k-1}v = uy^{n-k-1}xv = uyxv.$$

Since, for every $a \in N$, there is a positive integer m such that $a^m \in K,\ a \,\beta\, a^m$, we get that S is a β class. Hence S is medial.

Consider the case when $j > 1$. Let α be an equivalence of S such that $a \,\alpha\, b \Leftrightarrow \forall x \in K\ xa = xb$ and $ax = bx$.
It is obvious that $x \,\alpha\, y$ implies $x = y$ for every $x, y \in K$. We show that α is also a congruence. Let $y, a, b \in S$ be arbitrary elements. Assume $a \,\alpha\, b$ then

$$x(ay) = (xa)y = (xb)y = x(by)$$

and

$$(ay)x = a(yx) = b(yx) = (by)x$$

and so $ay \,\alpha\, by$. We can prove $ya \,\alpha\, yb$ in a similar way. Assume that N is not nilpotent. Let $a_1, a_2, \cdots, a_j \in S - K$ be such that $a_1 a_2 \ldots a_j \in S - K$. We prove that $a_1 a_2 \ldots a_j \,\alpha\, a_1 a_2 \ldots a_j^m$ where m is the least positive integer such that $a_j^m \in K$. Since $a_1 a_2 \ldots a_j \in S - K$ then $a_{i_1} a_{i_2} = a_{i_2} a_{i_1}$ for every $i_1, i_2 \in \{1, 2, \cdots, j\}$. Let $l^{'}$ and $k^{'}$ be such that $\sigma(l+1) = l^{'}$ and $\sigma(n-k) = k^{'}$. Then, for every $x \in K$,

$$a_1 a_2 \ldots a_j x = a_1 a_2 \ldots a_j x^{l^{'}-l-1} a_j x^{n-l^{'}} x = a_1 a_2 \ldots a_j a_j x^{n-l-1} x = a_1 a_2 \ldots a_j^2 x$$

and so

$$a_1 a_2 \ldots a_j x = a_1 a_2 \ldots a_j^m x.$$

Using also $a_{i_1}a_{i_2} = a_{i_2}a_{i_1}$ $(i_1, i_2 \in \{1, 2, \cdots, j\})$, we have

$$xa_1a_2 \dots a_j = xa_1a_2 \dots a_j^m$$

in a similar way. From these results we have

$$a_1a_2 \dots a_j \, \alpha \, a_1a_2 \dots a_j^m.$$

From (2) of Lemma 3 of [3], we get $\mid K \mid = 1$, and so S is commutative. □

Theorem 2 *If S is an ideal extension of a rectangular band K by a non trivial commutative nilpotent semigroup N, and S is permutable then S is commutative.*

Proof. Assume that a permutable semigroup S is an ideal extension of a rectangular band K by a non trivial commutative nilpotent semigroup N. By Lemma 2 of [4], N is a Δ-semigroup. Then, by Lemma 2 of [5], N is finite ciclyc. Let $N = \{a, a^2, \cdots, a^n\}$ where $a^n \in K$. Assume $a^n = (1, 1)$ and $K = L \times R$, where L is a leftzero, R is a rightzero semigroup. For every $(x_1, x_2) \in K$, $a(x_1, x_2) = (\phi_a(x_1, x_2), x_2)$ and $(x_1, x_2)a = (x_1, \psi_a(x_1, x_2))$.
If $(x_1, x_3) \in K$, then

$$a(x_1, x_3) = a(x_1, x_2)(x_2, x_3) = (\phi_a(x_1, x_2), x_2)(x_2, x_3) = (\phi_a(x_1, x_2), x_3)$$

and so $\phi_a(x_1, x_2) = \phi_a(x_1, x_3)$. So ϕ_a is independent from the second variable. Similarly, ψ_a is independent from the first variable. From these results we get that, for every $(x_1, x_2) \in K$, $a(x_1, x_2) = (\phi_a(x_1), x_2)$ and $(x_1, x_2)a = (x_1, \psi_a(x_2))$. We will prove that $\phi_a(x) = 1$ for every $x \in L$. It is obvious that $\underbrace{\phi_a \circ \ldots \circ \phi_a}_{\text{n}}(x) = \phi_a^n(x) = 1$ for every $x \in L$. Let j be such that $\phi_a^j(x) = \phi_a^{j+1}(x) = \ldots = \phi_a^n(x) = 1$ $\forall x \in L$. Assume that $\exists x \in L$ such that $\phi_a^{j-1}(x) \neq 1$. Let

$$P := \{x \in L : \phi_a^{j-1}(x) \neq 1\}$$

and

$$Q := \{(x_1, x_2) \in K : x_1 \in P\}.$$

It is clear that, for every $(x_1, x_2) \in K$, $a^r(x_1, x_2) \notin Q$, for all positive integers r, because $a^r(x_1, x_2) = (\phi_a^r(x_1), x_2)$ and $\phi_a^{j-1}(\phi_a^r(x_1)) = \phi_a^{j+r-1}(x_1) = 1$. Since $\phi_a(1) = 1$ then $Q \neq K$.
Let α be an equivalence on S such that, for arbitrary $x, y \in S$, $x \, \alpha \, y \Leftrightarrow x, y \in Q$ or $x, y \in S - Q$. We show that α is a congruence of S. Let $x, y, z \in S$ be arbitrary elements with $x \, \alpha \, y$. Assume that $x, y \in K$. It is clear that if $z \in K$ then $xz \, \alpha \, yz$ and $zx \, \alpha \, zy$. Assume $z = a^l \in S - K$. It is obvious that $xa^l \, \alpha \, ya^l$. Since $a^lx, a^ly \notin Q$ then $a^lx \, \alpha \, a^ly$ is obvious. Consider the case when $x = a^m \in S - K$ and $y \in K$ where m is a positive integer less than n. For every $z \in K$ $zx \, \alpha \, zy$, is obvious. Since $xz, y \notin Q$, we get that $xz \, \alpha \, yz$. Consider the case when $x, y \in S - K$, that is $x = a^p$ and $y = a^q$, where p, q are positive integers less than n. For all $z \in S$, $zx \, \alpha \, zy$ and $xz \, \alpha \, yz$ are obvious. So α is a congruence,

indeed. There are two α classes on S: Q and $S-Q$. It is clear that $a^n \in K-Q$ and so $Q \neq K$. From (2) of Lemma 3 of [3], K have to be union of α classes or subset of an α class. Since $(S-Q)\cap K \neq 0$, $(S-Q)\cap(S-K) \neq 0$ and $Q \subset K$, then Q have to be an empty set! Since $\mid Q \mid \geq \mid P \mid$, then $0 = \mid Q \mid \geq \mid P \mid \geq 0$. That means $\mid P \mid = 0$, that is $\phi_a(L) = \{1\}$. We can prove $\psi_a(R) = \{1\}$ in a similar way. Consequently, for every $x \in K$, $ax = a^2x = \ldots = a^nx$ and $xa = xa^2 = \ldots = xa^n$.
Let β be an equivalence of S such that $x\,\beta\,y \Leftrightarrow x = y$ or $x = a^l,\ y = a^k$ where l, k are positive integers. Let $x, y, z \in S$ be arbitrary elements with $x\,\beta\,y$. If $x = y$ then $xz\,\beta\,yz$ and $zx\,\beta\,zy$ are obivous. If $x \neq y$ then, with $x\,\beta\,y$, we get that $x = a^l$ and $y = a^k$, for some positive integers l, k. If $z = a^m$ then $a^m a^l\,\alpha\,a^m a^k$ and $a^l a^m\,\alpha\,a^k a^m$ are obvious. Let $x \in K$, then $xa^l = xa^n = xa^k$ and $a^l x = a^n x = a^k x$. Hence β is a congruence. It is clear that, for all $x, y \in K$, $x\,\beta\,y \Leftrightarrow x = y$ and, for all $x, y \in \{a, a^2, \cdots, a^n\}$, $x\,\beta\,y$. So the β classes of S are: $\{a, a^2, \cdots, a^n\} = N$ and for every $x \in S - N$, x is a β class. From (2) of Lemma 3 of [3] K have to be union of β classes, or subset of a β class. Since $\{a, a^2, \cdots, a^n\} \cap K = a^n$ and $S = \{a, a^2, \cdots, a^{n-1}\} \cup K$, then $K = a^n$. That is $\mid K \mid = 1$. And so S is commutative. □

Corollary 1 *Every permutable semigroup satisfying a non-trivial permutation identity is medial.*

Proof. From Theorem 2 of [4] and Theorem 1 and Theorem 2 of this paper it is obvious. □

References

[1] Clifford, A.H. and G.B. Preston, The Algebraic Theory of Semigroups, Amer. Math. Soc., Providence, R.I., I(1961), II(1967)

[2] Hamilton, H., Permutability of congruences on commutative semigroups, Semigroup Forum, 10(1975), 55-66

[3] Jiang, Z., LC-commutative permutable semigroups, Semigroup Forum, 52(1995), 191-196

[4] Nagy, A., Permutative semigroups satisfying a non-trivial permutation identity, Acta Sci. Math. (Szeged), 71(2005), 37-43

[5] Nagy, A. and P.R. Jones, Permutative semigroups whose congruences form a chain, Semigroup Forum (to appear)

Attila Deák, Department of Algebra, Institute of Mathematics, Budapest University of Technology and Economics; e-mail: ignotus@math.bme.hu